

ON DENSE ORBITS IN THE BOUNDARY OF A COXETER SYSTEM

TETSUYA HOSAKA

ABSTRACT. In this paper, we study the minimality of the boundary of a Coxeter system. We show that for a Coxeter system (W, S) if there exist a maximal spherical subset T of S and an element $s_0 \in S$ such that $m(s_0, t) \geq 3$ for each $t \in T$ and $m(s_0, t_0) = \infty$ for some $t_0 \in T$, then every orbit $W\alpha$ is dense in the boundary $\partial\Sigma(W, S)$ of the Coxeter system (W, S) , hence $\partial\Sigma(W, S)$ is minimal, where $m(s_0, t)$ is the order of $s_0 t$ in W .

1. INTRODUCTION AND PRELIMINARIES

The purpose of this paper is to study the minimality of the boundary of a Coxeter system. A *Coxeter group* is a group W having a presentation

$$\langle S \mid (st)^{m(s,t)} = 1 \text{ for } s, t \in S \rangle,$$

where S is a finite set and $m : S \times S \rightarrow \mathbb{N} \cup \{\infty\}$ is a function satisfying the following conditions:

- (1) $m(s, t) = m(t, s)$ for each $s, t \in S$,
- (2) $m(s, s) = 1$ for each $s \in S$, and
- (3) $m(s, t) \geq 2$ for each $s, t \in S$ such that $s \neq t$.

The pair (W, S) is called a *Coxeter system*. Let (W, S) be a Coxeter system. For a subset $T \subset S$, W_T is defined as the subgroup of W generated by T , and called a *parabolic subgroup*. If T is the empty set, then W_T is the trivial group. A subset $T \subset S$ is called a *spherical subset of S* , if the parabolic subgroup W_T is finite.

Every Coxeter system (W, S) determines a *Davis-Moussong complex* $\Sigma(W, S)$ which is a CAT(0) geodesic space ([4], [5], [6], [10]). Here the 1-skeleton of $\Sigma(W, S)$ is the Cayley graph of W with respect to S . The natural action of W on $\Sigma(W, S)$ is proper, cocompact and by isometry.

Date: February 1, 2005.

2000 Mathematics Subject Classification. 57M07, 20F65, 20F55.

Key words and phrases. boundaries of Coxeter groups.

Partly supported by the Grant-in-Aid for Young Scientists (B), The Ministry of Education, Culture, Sports, Science and Technology, Japan. (No. 15740029).

We can consider a certain fundamental domain $K(W, S)$ which is called a *chamber* of $\Sigma(W, S)$ such that $WK(W, S) = \Sigma(W, S)$ ([5], [6]). If W is infinite, then $\Sigma(W, S)$ is noncompact and $\Sigma(W, S)$ can be compactified by adding its ideal boundary $\partial\Sigma(W, S)$ ([2], [5, §4]). This boundary $\partial\Sigma(W, S)$ is called the *boundary of* (W, S) . We note that the natural action of W on $\Sigma(W, S)$ induces an action of W on $\partial\Sigma(W, S)$.

The following theorem was proved in [8].

Theorem 1.1. *Let (W, S) be a Coxeter system. Suppose that there exist a maximal spherical subset T of S and an element $s_0 \in S$ such that $m(s_0, t) \geq 3$ for each $t \in T$ and $m(s_0, t_0) = \infty$ for some $t_0 \in T$. Then $W\alpha$ is dense in $\partial\Sigma(W, S)$ for some $\alpha \in \partial\Sigma(W, S)$.*

Suppose that a group G acts on a compact metric space X by homeomorphisms. Then X is said to be *minimal*, if every orbit Gx is dense in X .

For a negatively curved group Γ and the boundary $\partial\Gamma$ of Γ , by an easy argument, we can show that $\Gamma\alpha$ is dense in $\partial\Gamma$ for each $\alpha \in \partial\Gamma$, that is, $\partial\Gamma$ is minimal.

In this paper, we prove the following theorem as an extension of Theorem 1.1.

Theorem 1.2. *Let (W, S) be a Coxeter system which satisfies the condition in Theorem 1.1. Then every orbit $W\alpha$ is dense in $\partial\Sigma(W, S)$, that is, $\partial\Sigma(W, S)$ is minimal.*

2. LEMMAS ON COXETER GROUPS AND THE DAVIS-MOUSSONG COMPLEXES

In this section, we recall and prove some lemmas for Coxeter groups and the Davis-Moussong complexes which are used later.

Definition 2.1. Let (W, S) be a Coxeter system and $w \in W$. A representation $w = s_1 \cdots s_l$ ($s_i \in S$) is said to be *reduced*, if $\ell(w) = l$, where $\ell(w)$ is the minimum length of word in S which represents w .

The following lemma is known.

Lemma 2.2 ([1], [3], [4], [6], [9]). *Let (W, S) be a Coxeter system.*

- (1) *Let $w \in W$ and let $w = s_1 \cdots s_l$ be a representation. If $\ell(w) < l$, then $w = s_1 \cdots \hat{s}_i \cdots \hat{s}_j \cdots s_l$ for some $1 \leq i < j \leq l$.*
- (2) *For each $w \in W$ and $s \in S$, $\ell(ws)$ equals either $\ell(w) + 1$ or $\ell(w) - 1$, and $\ell(sw)$ also equals either $\ell(w) + 1$ or $\ell(w) - 1$.*
- (3) *For each $w \in W$, $S(w)$ is a spherical subset of S , i.e., $W_{S(w)}$ is finite.*

- (4) For each $w \in W$ and each spherical subset T of S , there exists a unique element of longest length in $W_T w$. Here $v \in W_T w$ is the element of longest length in $W_T w$ if and only if $\ell(tv) < \ell(v)$ for any $t \in T$. Then $\ell(v) = \ell(vw^{-1}) + \ell(w)$.

Lemma 2.3. Let (W, S) be a Coxeter system, let $w \in W$, let $s \in S$ such that $\ell(ws) = \ell(w) + 1$, and let T be a spherical subset of S . By Lemma 2.2 (4), there exist unique elements $x, x' \in W_T$ such that xw and $x'ws$ are the elements of longest length in $W_T w$ and $W_T ws$ respectively. If $x = t_1 \dots t_m$ is a reduced representation, then either $x' = x$ or $x' = t_1 \dots \hat{t}_i \dots t_m$ for some $i \in \{1, \dots, m\}$. Hence $\ell(x') \leq \ell(x)$.

Proof. By Lemma 2.2 (2), either $\ell(xws) = \ell(xw) + 1$ or $\ell(xws) = \ell(xw) - 1$.

We first suppose that $\ell(xws) = \ell(xw) + 1$. Since xw is the element of longest length in $W_T w$, $\ell(twx) < \ell(xw)$ for any $t \in T$ by Lemma 2.2 (4). Then for each $t \in T$,

$$\ell(txws) \leq \ell(twx) + 1 < \ell(xw) + 1 = \ell(xws).$$

Hence $\ell(txws) < \ell(xws)$ for any $t \in T$. Thus xws is the element of longest length in $W_T ws$, i.e., $x' = x$.

Next we suppose that $\ell(xws) = \ell(xw) - 1$. Since $\ell(ws) = \ell(w) + 1$ and $\ell(xw) = \ell(x) + \ell(w)$,

$$xws = (t_1 \dots t_m)ws = (t_1 \dots \hat{t}_i \dots t_m)w$$

for some $i \in \{1, \dots, m\}$ by Lemma 2.2 (1). Now xw is the element of longest length in $W_T w$. Here

$$W_T w = W_T(t_1 \dots \hat{t}_i \dots t_m)w = W_T xws = W_T ws.$$

Hence xw is the element of longest length in $W_T ws$. Since $xw = (t_1 \dots \hat{t}_i \dots t_m)ws$, we obtain $x' = t_1 \dots \hat{t}_i \dots t_m$. $\square$

Definition 2.4. Let (W, S) be a Coxeter system. For each $w \in W$, we define $S(w) = \{s \in S \mid \ell(ws) < \ell(w)\}$. For a subset $T \subset S$, we also define $W^T = \{w \in W \mid S(w) = T\}$.

The following lemma was proved in [8].

Lemma 2.5 ([8, Lemma 2.5]). Let (W, S) be a Coxeter system, $w \in W$ and $s_0 \in S$. Suppose that $m(s_0, t) \geq 3$ for each $t \in S(w)$ and that $m(s_0, t_0) = \infty$ for some $t_0 \in S(w)$. Then $ws_0 \in W^{\{s_0\}}$.

We can obtain the following lemma by the same argument as the proof of [7, Lemma 4.2].

Lemma 2.6. *Let (W, S) be a Coxeter system and let $\alpha \in \partial\Sigma(W, S)$. Then there exists a sequence $\{s_i\} \subset S$ such that $s_1 \cdots s_i$ is reduced and $d(s_1 \cdots s_i, \text{Im } \xi_\alpha) \leq N$ for each $i \in \mathbb{N}$, where N is the diameter of $K(W, S)$ in $\Sigma(W, S)$ and ξ_α is the geodesic ray in $\Sigma(W, S)$ such that $\xi_\alpha(0) = 1$ and $\xi_\alpha(\infty) = \alpha$.*

3. PROOF OF THE MAIN THEOREM

Using the lemmas above, we prove Theorem 1.2.

Proof of Theorem 1.2. Let (W, S) be a Coxeter system. Suppose that there exist a maximal spherical subset T of S and an element $s_0 \in S$ such that $m(s_0, t) \geq 3$ for each $t \in T$ and $m(s_0, t_0) = \infty$ for some $t_0 \in T$. Let $\alpha \in \partial\Sigma(W, S)$. By Lemma 2.6, there exists a sequence $\{s_i\} \subset S$ such that $s_1 \cdots s_i$ is reduced and $d(s_1 \cdots s_i, \text{Im } \xi_\alpha) \leq N$ for each $i \in \mathbb{N}$, where N is the diameter of $K(W, S)$ in $\Sigma(W, S)$ and ξ_α is the geodesic ray such that $\xi_\alpha(0) = 1$ and $\xi_\alpha(\infty) = \alpha$. Let $w_i = s_1 \cdots s_i$. For each i , there exists a unique element $x_i \in W_T$ such that $x_i w_i$ is the element of longest length in $W_T w_i$ by Lemma 2.2 (4). Now $w_{i+1} = w_i s_{i+1}$ and $\ell(w_{i+1}) = \ell(w_i) + 1$. By Lemma 2.3, $\ell(x_{i+1}) \leq \ell(x_i)$ for every i . Hence there exists a number n such that $\ell(x_i) = \ell(x_{i+1})$ for each $i \geq n$. Then $x_i = x_{i+1}$ for every $i \geq n$ by Lemma 2.3. Let $x = x_n$. Then $x w_i$ is the element of longest length in $W_T w_i$ for each $i \geq n$. Since $\ell(t x w_i) < \ell(x w_i)$ for any $t \in T$, $T \subset S((x w_i)^{-1})$. Here $S((x w_i)^{-1})$ is a spherical subset of S by Lemma 2.2 (3) and T is a maximal spherical subset of S . Hence $S((x w_i)^{-1}) = T$ for each $i \geq n$. By Lemma 2.5, $(s_0 x w_i)^{-1} \in W^{\{s_0\}}$ and $(t_0 s_0 x w_i)^{-1} \in W^{\{t_0\}}$ for each $i \geq n$. Hence $(W^{\{t_0\}})^{-1}$ contains the sequence $\{t_0 s_0 x w_i\}_{i \geq n}$ which converges to $t_0 s_0 x \alpha$. By the proof of [8, Theorem 4.1], $W t_0 s_0 x \alpha$ is dense in $\partial\Sigma(W, S)$. Here $W t_0 s_0 x \alpha = W \alpha$. Hence $W \alpha$ is a dense subset of $\partial\Sigma(W, S)$. Thus every orbit $W \alpha$ is dense in $\partial\Sigma(W, S)$, that is, $\partial\Sigma(W, S)$ is minimal. $\square$

REFERENCES

- [1] N. Bourbaki, *Groupes et Algèbres de Lie*, Chapters IV-VI, Masson, Paris, 1981.
- [2] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Springer-Verlag, Berlin, 1999.
- [3] K. S. Brown, *Buildings*, Springer-Verlag, 1980.
- [4] M. W. Davis, *Groups generated by reflections and aspherical manifolds not covered by Euclidean space*, Ann. of Math. **117** (1983), 293–324.
- [5] ———, *Nonpositive curvature and reflection groups*, in Handbook of geometric topology (Edited by R. J. Daverman and R. B. Sher), pp. 373–422, North-Holland, Amsterdam, 2002.

- [6] ———, *The cohomology of a Coxeter group with group ring coefficients*, Duke Math. J. **91** (no.2) (1998), 297–314.
- [7] T. Hosaka, *Parabolic subgroups of finite index in Coxeter groups*, J. Pure Appl. Algebra **169** (2002), 215–227.
- [8] ———, *Dense subsets of the boundary of a Coxeter system*, Proc. Amer. Math. Soc. **132** (2004), 3441–3448.
- [9] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge University Press, 1990.
- [10] G. Moussong, *Hyperbolic Coxeter groups*, Ph.D. thesis, The Ohio State University, 1988.

DEPARTMENT OF MATHEMATICS, UTSUNOMIYA UNIVERSITY, UTSUNOMIYA,
 321-8505, JAPAN
E-mail address: hosaka@cc.utsunomiya-u.ac.jp